

Midterm Formula Sheet

Constants:

$$h = 6.63 \times 10^{-34} \text{ J-s} = 4.13 \times 10^{-15} \text{ eV-s}$$

$$\hbar = h / 2\pi = 1.05 \times 10^{-34} \text{ J-s} = 6.58 \times 10^{-16} \text{ eV-s}$$

$$k = 1.38 \times 10^{-23} \text{ J/K} = 8.62 \times 10^{-5} \text{ eV/K},$$

$$m_0 = 9.11 \times 10^{-31} \text{ kg},$$

$$kT/q = 0.0259 \text{ V at } 300 \text{ K}$$

$$c = 3 \times 10^{10} \text{ cm/s}$$

$$q = 1.60 \times 10^{-19} \text{ C}$$

Equations:

$$J = \frac{I}{A} = \sigma E = (q\mu_n n + q\mu_p p)E, \quad R = \rho \frac{L}{S}, \quad \sigma = \frac{1}{\rho}, \quad \sigma = q\mu_n n + q\mu_p p$$

$$\text{Photoelectric experiment: } K.E._{\text{max}} = h\nu - \Phi$$

$$\text{Time-dependent: } -\frac{\hbar^2}{2m} \nabla^2 \psi(\vec{r}, t) + U(\vec{r}, t)\psi(\vec{r}, t) = j\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t}$$

$$\text{Time-independent: } -\frac{\hbar^2}{2m} \nabla^2 \psi(\vec{r}) + U(\vec{r})\psi(\vec{r}) = E\psi(\vec{r})$$

$$\text{Wave number: } k = \frac{\sqrt{2m(E - U_0)}}{\hbar} \text{ for } E > U_0 \text{ and } k = \frac{\sqrt{2m(U_0 - E)}}{\hbar} \text{ for } E < U_0$$

$$\text{Momentum: } p = m^* v = \hbar k = \frac{h}{\lambda}, \text{ operator: } \frac{\hbar}{j} \frac{\partial}{\partial x}; \text{ Energy: } E = h\nu = \hbar\omega, \text{ operator: } -\frac{\hbar}{j} \frac{\partial}{\partial t}$$

$$\text{Kinetic energy: } K.E. = \frac{m^* v^2}{2} = \frac{\hbar^2 k^2}{2m^*}, \text{ operator: } -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2}$$

$$k = \frac{2\pi}{\lambda}, \quad \omega = 2\pi\nu,$$

$$\text{Hydrogen atom: } E(n) = \frac{-13.6}{n^2} \text{ eV}$$

$$\text{Mobility: } \mu_d = \frac{q\tau}{m}, \quad v_d = \mu_d E$$

$$\text{Fermi-Dirac distribution: } f(E) = \frac{1}{1 + e^{(E - E_F)/kT}} \cong e^{-(E - E_F)/kT} \text{ for } E - E_F \gg kT$$

$$n = N_C \exp\left(\frac{E_F - E_C}{kT}\right) = n_i \exp\left(\frac{E_F - E_{Fi}}{kT}\right), \quad p = N_V \exp\left(\frac{E_V - E_F}{kT}\right) = n_i \exp\left(\frac{E_{Fi} - E_F}{kT}\right)$$

$$N_C = 2 \left(\frac{2\pi m_n^* kT}{h^2} \right)^{3/2}, \quad N_V = 2 \left(\frac{2\pi m_p^* kT}{h^2} \right)^{3/2}, \quad n_i = B \exp\left(\frac{-E_g}{2kT}\right) = \sqrt{N_C N_V} \exp\left(\frac{-E_g}{2kT}\right)$$

$$\text{Equilibrium: } n_0 p_0 = n_i^2$$

$$\text{Charge neutrality: } n_0 + N_A = p_0 + N_D$$

$$\text{Recombination: } r(T) = \alpha_r n p = \alpha_r n_0 p_0 + \alpha_r n_0 \hat{p} + \alpha_r \hat{n} p_0$$

$$\hat{n}(t) = \hat{n}(0) \exp(-t/\tau_n),$$

$$\hat{p}(t) = \hat{p}(0) \exp(-t/\tau_p)$$